



Cell Detachment Solution

Catalog #: ADF001

Size: 50 ml

Storage: Store at -20°C until ready for use and 2-8°C for short term

Format: Liquid

GENERAL INFORMATION

The Cell Detachment Solution is an effective alternative to Trypsin/EDTA for in vitro cultured cells, containing proteolytic and collagenolytic enzymes for the usage of detachment of cells from standard tissue culture plates or T-flasks. Cell Detachment Solution contains protease and collagenase activities in an isotonic, phosphate buffer solution with EDTA to successfully to detach primary cells and cell lines.

Benefits of Cell Detachment Solution:

- Maximum cell viability
- No need to wash detached cells
- Gentle and rapid detachment
- Effective on a wide range of cells
- No mammalian or bacterial by-products

Shipping and Storage

Product can be shipped with an ice pack or dry ice. Cell Detachment Solution can be stored at 2°C-8°C for up two months or -20°C for long term storage. Alternatively, if the product is received frozen, wait to thaw until ready for use.

Protocol

1. Thaw Cell Detachment Solution at room temperature
2. Remove cell culture media
3. Rinse culture plates or T-flasks with 1XPBS
4. Dispose of 1XPBS by aspiration
5. Add an adequate volume of Cell Detachment Solution to cover cells. Use minimum of 1.0 ml per 75 cm² surface area. Rock container back and forth to cover surface area with solution
6. Allow cells to detach at room temperature or 37°C Incubator for approximately 5-10 minutes or until the cells are detached
7. Monitor cell detachment. Strike flask against palm of the hand to detach cells if needed
8. After cells are detached, add growth media at a volume equal to the original enzyme solution
9. Gently pipette the cells suspension up and down five times with the culture plates or T-flask in order to flush out the remaining cells from the culture plate or T-flask
10. Count cells, passage, or process as needed
11. Cells are ready to be used for reseeding or experimental analysis

Note: Proper precautions must be taken to avoid inadvertent exposure. Always wear gloves and safety glasses when working these materials. Never mouth pipette. We recommend following the universal procedures for handling products of human origin as the minimum precaution against contamination.

FOR RESEARCH USE ONLY

NEUROMICS REAGENTS ARE FOR IN VITRO AND CERTAIN NON-HUMAN IN VIVO EXPERIMENTAL USE ONLY AND NOT INTENDED FOR USE IN ANY HUMAN CLINICAL INVESTIGATION, DIAGNOSIS, PROGNOSIS, OR TREATMENT. THE ABOVE ANALYSES ARE MERELY TYPICAL GUIDES. THEY ARE NOT TO BE CONSTRUED AS BEING SPECIFICATIONS. ALL OF THE ABOVE INFORMATION IS, TO THE BEST OF OUR KNOWLEDGE, TRUE AND ACCURATE. HOWEVER, SINCE THE CONDITIONS OF USE ARE BEYOND OUR CONTROL, ALL RECOMMENDATIONS OR SUGGESTIONS ARE MADE WITHOUT GUARANTEE, EXPRESS OR IMPLIED, ON OUR PART. WE DISCLAIM ALL LIABILITY IN CONNECTION WITH THE USE OF THE INFORMATION CONTAINED HEREIN OR OTHERWISE, AND ALL SUCH RISKS ARE ASSUMED BY THE USER. WE FURTHER EXPRESSLY DISCLAIM ALL WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE. v1-09809

www.neuromics.com

Neuromics Antibodies • 5325 West 74th Street, Suite 8 • Edina, MN 55439
phone 866-350-1500 • fax 612-677-3976 • e-mail pshuster@neuromics.com