



Catalog Number:	RA18021	Host:	Rabbit
Product Type:	Affinity purified antibody	Species Reactivity:	Rat, Mouse, Human, Primate
Immunogen Sequence:	Peptide derived from the carboxy-terminal sequence of human alpha-synuclein.	Format:	Liquid in 10mM sodium HEPES (pH7.5), 150mM NaCl, 100ug BSA and 50% glycerol
Applications:	Western blotting 1:1000 Immunoprecipitation 1:50 Dilutions listed only as a recommendation. Optimal dilution should be determined by investigator.		
Storage:	Store at -20°C. Do not aliquot.		

Application Notes

alpha-Synuclein antibody detects the alpha isoform of synuclein. The antibody may cross-react with beta-synuclein.

Western Blot Protocol

Sample Preparation

1. Treat cells by adding fresh media containing regulator for desired time.
2. Aspirate media from cultures; wash cells with 1X PBS; aspirate.
3. Lyse cells by adding 1X SDS sample buffer (100 μ l per well of 6-well plate or 500 μ l per plate of 10 cm plate). Immediately scrape cells from plate and transfer the extract to a microcentrifuge tube. Keep on ice.
4. Sonicate for 10–15 seconds to shear DNA and reduce sample viscosity.
5. Heat a 20 μ l sample to 95–100°C for 5 minutes; cool on ice.
6. Microcentrifuge for 5 minutes.
7. Load 20 μ l onto SDS-PAGE gel (10 cm x 10 cm).
8. Electrotransfer to nitrocellulose (or PVDF) membrane.

Membrane Blocking and Antibody Incubations

Note: Volumes for 10 cm x 10 cm (100 cm²) membrane; for different sized membranes, adjust vol. accordingly.

1. (Optional) After transfer, wash nitrocellulose membrane with 25 ml TBS for 5 minutes at room temperature.
2. Incubate membrane in 25 ml of blocking buffer for 1 hour at room temperature.
3. Wash three times for 5 minutes each with 15 ml of TBS/T.
4. Incubate membrane and primary antibody (at the appropriate dilution) in 10 ml primary antibody dilution buffer with gentle agitation overnight at 4°C.
5. Wash three times for 5 minutes each with 15 ml of TBS/T.
6. Incubate membrane with HRP-conjugated secondary antibody (1:2000) in 10 ml of blocking buffer with gentle agitation for 1 hour at room temperature.
7. Wash three times for 5 minutes each with 15 ml of TBS/T.
8. Process membranes using enhanced chemiluminescence.

FOR RESEARCH USE ONLY

Data and Protocol Provided Courtesy of Cell Signaling Technology, Inc.

NEUROMICS REAGENTS ARE FOR IN VITRO AND CERTAIN NON-HUMAN IN VIVO EXPERIMENTAL USE ONLY AND NOT INTENDED FOR USE IN ANY HUMAN CLINICAL INVESTIGATION, DIAGNOSIS, PROGNOSIS, OR TREATMENT. THE ABOVE ANALYSES ARE MERELY TYPICAL GUIDES. THEY ARE NOT TO BE CONSTRUED AS BEING SPECIFICATIONS. ALL OF THE ABOVE INFORMATION IS, TO THE BEST OF OUR KNOWLEDGE, TRUE AND ACCURATE. HOWEVER, SINCE THE CONDITIONS OF USE ARE BEYOND OUR CONTROL, ALL RECOMMENDATIONS OR SUGGESTIONS ARE MADE WITHOUT GUARANTEE, EXPRESS OR IMPLIED, ON OUR PART. WE DISCLAIM ALL LIABILITY IN CONNECTION WITH THE USE OF THE INFORMATION CONTAINED HEREIN OR OTHERWISE, AND ALL SUCH RISKS ARE ASSUMED BY THE USER. WE FURTHER EXPRESSLY DISCLAIM ALL WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE.

www.neuromics.com

Neuromics Antibodies • 5325 West 74th Street, Suite 8 • Edina, MN 55439
phone 866-350-1500 • fax 612-677-3976 • e-mail pshuster@neuromics1.com

10/04v1

Immunoprecipitation Protocol**Sample Preparation**

1. Aspirate media. Treat cells by adding fresh media containing regulator for desired time.
2. To harvest cells under non-denaturing conditions, remove media and rinse cells once with ice-cold PBS.
3. Remove PBS and add 0.5 ml 1X ice-cold cell lysis buffer plus 1 mM PMSF to each plate (10 cm) and incubate the plate on ice for 5 minutes.
4. Scrape cells off the plate and transfer to microcentrifuge tubes. Keep on ice.
5. Sonicate on ice four times for 5 seconds each.
6. Microcentrifuge for 10 minutes at 4°C, and transfer the supernatant to a new tube. The supernatant is the cell lysate. If necessary, lysate can be stored at -80°C.

Immunoprecipitation

1. Take 200 µl cell lysate and add primary antibody; incubate with gentle rocking overnight at 4°C.
2. Add protein A agarose beads (20 µl of 50% bead slurry). Incubate with gentle rocking for 1–3 hrs at 4°C.
3. Microcentrifuge for 30 seconds at 4°C. Wash pellet five times with 500 µl of 1X cell lysis buffer. Keep on ice during washes.
4. Resuspend the pellet with 20 µl 3X SDS sample buffer. Vortex, then microcentrifuge for 30 seconds.
5. Heat the sample to 95–100°C for 2–5 minutes.
6. Load the sample (15–30 µl) on SDS-PAGE gel (12–15%).
7. Analyze sample by Western blotting (see Western Immunoblotting Protocol).

Solutions and Reagents for Western Blot and Immunoprecipitation

Note: Prepare solutions with Milli-Q or equivalently purified water.

1X Phosphate Buffered Saline (PBS)**1X SDS Sample Buffer:**

62.5 mM Tris-HCl (pH 6.8 at 25°C), 2% w/v SDS, 10% glycerol, 50 mM DTT, 0.01% w/v bromophenol blue or phenol red

3X SDS Sample Buffer:

187.5 mM Tris-HCl (pH 6.8 at 25°C), 6% w/v SDS, 30% glycerol, 150 mM DTT, 0.03% w/v bromophenol blue

1X Cell Lysis Buffer:

20 mM Tris (pH 7.5), 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1% Triton X-100, 2.5 mM Sodium pyrophosphate, 1 mM β-Glycerolphosphate, 1 mM Na3VO4, 1 µg/ml Leupeptin. *Note: Addition of PMSF before use is suggested.*

Transfer Buffer:

25 mM Tris base, 0.2 M glycine, 20% methanol (pH 8.5)

10X Tris Buffered Saline (TBS):

To prepare 1 liter of 10X TBS: 24.2 g Tris base, 80 g NaCl; adjust pH to 7.6 with HCl (use at 1X).

Nonfat Dry Milk (weight to volume [w/v])**Blocking Buffer:**

1X TBS, 0.1% Tween-20 with 5% w/v nonfat dry milk; for 150 ml, add 15 ml 10X TBS to 135 ml water, mix. Add 7.5 g nonfat dry milk and mix well. While stirring, add 0.15 ml Tween-20 (100%).

Wash Buffer:

1X TBS, 0.1% Tween-20 (TBS/T)

Bovine Serum Albumin (BSA)**Primary Antibody Dilution Buffer:**

1X TBS, 0.1% Tween-20 with 5% BSA; for 20 ml, add 2 ml 10X TBS to 18 ml water, mix. Add 1.0 g BSA and mix well. While stirring, add 20 µl Tween-20 (100%).

Protein A Agarose Beads:

Add 5 ml of 1X PBS to 1.5 g of protein A agarose beads. Shake 2 hours at 4°C; spin down. Wash pellet twice with PBS. Resuspend beads in 1 volume of PBS. (Can be stored for 2 weeks at 4°C).

FOR RESEARCH USE ONLY**Data and Protocol Provided Courtesy of Cell Signaling Technology, Inc.**

NEUROMICS REAGENTS ARE FOR IN VITRO AND CERTAIN NON-HUMAN IN VIVO EXPERIMENTAL USE ONLY AND NOT INTENDED FOR USE IN ANY HUMAN CLINICAL INVESTIGATION, DIAGNOSIS, PROGNOSIS, OR TREATMENT. THE ABOVE ANALYSES ARE MERELY TYPICAL GUIDES. THEY ARE NOT TO BE CONSTRUED AS BEING SPECIFICATIONS. ALL OF THE ABOVE INFORMATION IS, TO THE BEST OF OUR KNOWLEDGE, TRUE AND ACCURATE. HOWEVER, SINCE THE CONDITIONS OF USE ARE BEYOND OUR CONTROL, ALL RECOMMENDATIONS OR SUGGESTIONS ARE MADE WITHOUT GUARANTEE, EXPRESS OR IMPLIED, ON OUR PART. WE DISCLAIM ALL LIABILITY IN CONNECTION WITH THE USE OF THE INFORMATION CONTAINED HEREIN OR OTHERWISE, AND ALL SUCH RISKS ARE ASSUMED BY THE USER. WE FURTHER EXPRESSLY DISCLAIM ALL WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE.

www.neuromics.com

Neuromics Antibodies • 5325 West 74th Street, Suite 8 • Edina, MN 55439
phone 866-350-1500 • fax 612-677-3976 • e-mail pshuster@neuromics1.com

10/04v1