



Catalog Number:	RA24428	Host:	Rabbit
Product Type:	Whole Serum	Species Reactivity:	Human, Monkey, Mouse, Pigeon, Rat
Immunogen Sequence:	Parvalbumin purified from rat muscle	Format:	100ul Lyophilized, \leq 0.09% sodium azide
Applications:	Immunohistochemistry: 1/5,000–1/8,000 in PBS/0.3% Triton X-100 - Bn/AV-HRP		

Storage and Preparation: Storage: Dilute with phosphate buffer or Tris buffer at dilutions no higher than 1/10, aliquot and freeze at -15° C or lower. Antibody can be stored for up to six months if handled as described above.

It is strongly recommended that the customer perform a primary antibody dilution series using our dilution recommendations as a guideline. Note that a change in the fixation or buffering system as used in our protocol may change the configuration of the protein and, therefore, may alter the reactivity with the tissue tested.

Application Notes for Immunohistochemistry

Tissue: Rat thalamus, hippocampus and cortex

Perfusion Fixation • Fixation: 4% paraformaldehyde in 0.1M phosphate buffer, pH 7.4; 500 mL over 20 min. • Post Fixation: 1.5 hour at 4°C in 4% paraformaldehyde in 0.1M phosphate buffer, pH 7.4. • Note: Paraformaldehyde is a necessary component of fixation for this antiserum. If needed for other applications, glutaraldehyde may be used at low levels (0.1–0.3%) in conjunction with paraformaldehyde.

Sections 10 μ m cryostat or 50 μ m vibratome

Tissue Incubation 18–24 hours at 2°–8°C.

Detection System Use IF or Bn-AV/HRP reagents at dilutions recommended by the manufacturers.

Image: IHC image of the rat hippocampus staining for parvalbumin. The tissue was fixed with 4% formaldehyde in phosphate buffer, before being removed and prepared for vibratome sectioning. Floating sections were incubated at RT in 10% goat serum in PBS, before standard IHC procedure. Primary antibody was incubated at 1:5000 for 48 hours, goat anti-rabbit secondary was subsequently added for 1 hour after washing with PBS. Light microscopy staining was achieved with standard biotin-streptavidin/HRP procedure and DAB chromogen.



FOR RESEARCH USE ONLY

NEUROMICS' REAGENTS ARE FOR IN VITRO AND CERTAIN NON-HUMAN IN VIVO EXPERIMENTAL USE ONLY AND NOT INTENDED FOR USE IN ANY HUMAN CLINICAL INVESTIGATION, DIAGNOSIS, PROGNOSIS, OR TREATMENT. THE ABOVE ANALYSES ARE MERELY TYPICAL GUIDES. THEY ARE NOT TO BE CONSTRUED AS BEING SPECIFICATIONS. ALL OF THE ABOVE INFORMATION IS, TO THE BEST OF OUR KNOWLEDGE, TRUE AND ACCURATE. HOWEVER, SINCE THE CONDITIONS OF USE ARE BEYOND OUR CONTROL, ALL RECOMMENDATIONS OR SUGGESTIONS ARE MADE WITHOUT GUARANTEE, EXPRESS OR IMPLIED, ON OUR PART. WE DISCLAIM ALL LIABILITY IN CONNECTION WITH THE USE OF THE INFORMATION CONTAINED HEREIN OR OTHERWISE, AND ALL SUCH RISKS ARE ASSUMED BY THE USER. WE FURTHER EXPRESSLY DISCLAIM ALL WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE.-V2/08/2012