

NEUROMICS



ISOKine™ EFGF

Datasheet

Catalog Number: PR80002

Product Type: Recombinant Protein

Source: Produced in the endosperm tissue of barley grain (*Hordeum vulgare*), that exhibits up to 50 times less protease activity than *E.coli* or mammalian cells. Barley seed is void of any human or animal viral contaminants that could jeopardize your cell culture.

Description/Molecular Mass: Recombinant human EGF contains 54 amino acids and a 16 a.a. histidine-based tag for a total length of 70 a.a. and has a predicted molecular mass of 8.5 kDa including his-tag. The recombinant protein migrates with an apparent molecular mass of 11 kDa in SDS-PAGE.

Activity: Each batch of ISOKine™ growth factor is tested for bioactivity and verified to have comparable activity to a commercial source. Bioactivity of ISOKine™ recombinant human EGF is assayed by measuring its dose dependent effect on proliferation of 3T3 cells. The ED50 for this effect using EGF is typically < 0.1 ng/ml, corresponding to a specific activity > 10 x10⁶ U/mg. Optimal concentration should be determined for specific applications and cell lines.

Endotoxin Level: Endotoxin level is less than 0.005ng per µg of ISOKine™ product (0.05EU/µg).

MAT Assay: Purified ISOKine™ product carries no pyrogenic or pro-inflammatory contaminants, as assayed with monocyte activation test using Human 10-plex Cytokine Assay measuring IL-6, TNF-alpha and IL-1beta induction.*** Ref. The Blood Bank, University Hospital of Iceland, Reykjavik, Iceland.

Purity: Greater than 95% by SDS-PAGE gel analysis

Format: Lyophilized . PBS, pH 7.2, sterile filtered.

Reconstitution: Always centrifuge the vial before opening. It is recommended to reconstitute the lyophilized protein in sterile water to a concentration of no less than 100 µg/ml. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

Storage: 12 months from date of receipt, -20 to 70°C as supplied.
1 month, 2 to 8 °C under sterile conditions after reconstitution.
3 months, - 20 to 70 °C under sterile conditions after reconstitution.
Avoid multiple freeze-thaw cycles.

FOR RESEARCH USE ONLY

NEUROMICS' REAGENTS ARE FOR IN VITRO AND CERTAIN NON-HUMAN IN VIVO EXPERIMENTAL USE ONLY AND NOT INTENDED FOR USE IN ANY HUMAN CLINICAL INVESTIGATION, DIAGNOSIS, PROGNOSIS, OR TREATMENT. THE ABOVE ANALYSES ARE MERELY TYPICAL GUIDES. THEY ARE NOT TO BE CONSTRUED AS BEING SPECIFICATIONS. ALL OF THE ABOVE INFORMATION IS, TO THE BEST OF OUR KNOWLEDGE, TRUE AND ACCURATE. HOWEVER, SINCE THE CONDITIONS OF USE ARE BEYOND OUR CONTROL, ALL RECOMMENDATIONS OR SUGGESTIONS ARE MADE WITHOUT GUARANTEE, EXPRESS OR IMPLIED, ON OUR PART. WE DISCLAIM ALL LIABILITY IN CONNECTION WITH THE USE OF THE INFORMATION CONTAINED HEREIN OR OTHERWISE, AND ALL SUCH RISKS ARE ASSUMED BY THE USER. WE FURTHER EXPRESSLY DISCLAIM ALL WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE.-V1-03/2014

www.neuromics.com

Neuromics • 5325 West 74th Street, Suite 8 • Edina, MN 55439
phone 866-350-1500 • fax 612-677-3976 • e-mail pshuster@neuromics.com