

P2Y2 Data Sheet

Catalog Number: P10139 Format: Liquid-50 ul-1mgs/ml

control

Application Notes

Basic Protocol for Absorption Control:

- Make 2 vials of your primary antibody at the working concentration specified on the data sheet or at the best concentration determined by you empirically. 1 vial is for normally processing with your primary antibody, the other vial is for use with the Blocking Peptide.
- Add the Blocking Peptide at the recommended dilution (see below) into 1 of the vials and incubate at room temperature for 1 hour.
- 3) Process your tissue/samples as normal with both antibody solutions.

Quick Calculation for Absorption Controls:

When we perform Absoption Controls, we use the peptide at a final concentration of ~10uM (1 x 10⁻⁵) or 1uM (1 x 10⁻⁶ M). Because most peptides are between 15-20 amino acids with a molecular weight of ~2000 Da, we supply most of our peptide at a concentration of 2mgs/ml. This allows for a simple dilution of 1:100 (~10 uM) or 1:1000 (~1uM) in the working antibody Solution.

FOR RESEARCH USE ONLY

NEUROMICS ANTIBODIES REAGENTS ARE FOR IN VITRO AND CERTAIN NON-HUMAN IN VIVO EXPERIMENTAL USE ONLY AND NOT INTENDED FOR USE IN ANY HUMAN CLINICAL INVESTIGATION, DIAGNOSIS, PROGNOSIS, OR TREATMENT. THE ABOVE ANALYSES ARE MERELY TYPICAL GUIDES. THEY ARE NOT TO BE CONSTRUED AS BEING SPECIFICATIONS. ALL OF THE ABOVE INFORMATION IS, TO THE BEST OF OUR KNOWLEDGE, TRUE AND ACCURATE. HOWEVER, SINCE THE CONDITIONS OF USE ARE BEYOND OUR CONTROL, ALL RECOMMENDATIONS OR SUGGESTIONS ARE MADE WITHOUT GUARANTEE, EXPRESS OR IMPLIED, ON OUR PART. WE DISCLAIM ALL LIABILITY IN CONNECTION WITH THE USE OF THE INFORMATION CONTAINED HEREIN OR OTHERWISE, AND ALL SUCH RISKS ARE ASSUMED BY THE USER. WE FURTHER EXPRESSLY DISCLAIM ALL WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE.