

# RFP Expressing Human Brain Glioblastoma Cells (LN-229)

Catalog #: TR06-RFP Cell #: >5x10<sup>5</sup> cells

Storage: Liquid Nitrogen until ready for culture. Product Format: Frozen Vial

While Culturing keep in 37°C CO<sub>2</sub> incubator

### **GENERAL INFORMATION**

Human Brain Glioblastoma Cells (LN-229) were derived from a Human Brain Glioblastoma of a 19-year old Caucasian female. RFP-Human Brain Glioblastoma Cells (LN-229) are selected from the Human Brain Glioblastoma Cells (LN-229) and transfected with RFP expressing lentiviruses resistant to puromycin. Cells are supplied in frozen vials with more than 5 x 10<sup>5</sup> cell/vial. Universal Full Growth Medium (TM001) is recommended to culture the cells.

Product is for Research use only.

Frozen Vials are shipped in a Dry Ice Package.

#### **CHARACTERIZATION OF THE CELLS**

Human Brain Glioblastoma cells (LN-229) are tested negative for HIV-1, HBV, HCV, and mycoplasma.

#### HANDLING OF ARRIVING CELLS

When you receive the dry ice package with cells in frozen vials, transfer the frozen vials of cells into a -80°C freezer for short period storage or a liquid nitrogen tank for long-term storage.

#### PROTOCOL FOR THAWING THE CELLS AND SUBCULTURE

**Note:** If you have any questions or need clarification regarding the protocol for culturing these cells, please reach out to Dr. Jensen Auguste at (978) 608-1766 with your questions before beginning.

- Pre-coating of T25 flasks-Add 2 ml each Universal Coating Solution (AC002) into a T25 flask to cover the
  whole surface of the flask, 5 mins later, dispose the excessive coating solution by aspiration and the flask is
  ready to be used (although solution containing other extracellular matrix, i.e. gelatin, collagen, and
  fibronectin, can be used, make sure to optimize the conditions in advance.
- 2. Thaw the frozen cell vial in a 37°C water bath first, and then transfer the cells into the pre-coated T25 flask with 10 ml of Universal Full Growth Medium (TM001), cells usually become confluent with 5-7 days.
- 3. To passage the cells, rinse the cells in a T25 flask with 5 ml HBSS (Room Temperature) twice; then add 2 ml Universal Detachment Solution (RT) (AD002) into one T25 flask; gently dispose the excessive Universal Detachment Solution within 20 seconds by aspiration.
- 4. Leave the T25 flask with the cells at RT or 37°C for 1 min (most cells usually will detach from the surface within 1-2 mins) or monitor the cells under a microscope until most of cells become rounded up, and then gently tap the flask against the bench surface, and the cells will move on the surface of the flask when monitoring under microscope.

#### **FUK KESEAKUTI USE UNLI**

NEUROMICS REAGENTS ARE FOR IN VITRO AND CERTAIN NON-HUMAN IN VIVO EXPERIMENTAL USE ONLY AND NOT INTENDED FOR USE IN ANY HUMAN CLINICAL INVESTIGATION, DIAGNOSIS, PROGNOSIS, OR TREATMENT. THE ABOVE ANALYSES ARE MERELY TYPICAL GUIDES. THEY ARE NOT TO BE CONSTRUED AS BEING SPECIFICATIONS. ALL OF THE ABOVE INFORMATION IS, TO THE BEST OF OUR KNOWLEDGE, TRUE AND ACCURATE. HOWEVER, SINCE THE CONDITIONS OF USE ARE BEYOND OUR CONTROL, ALL RECOMMENDATIONS OR SUGGESTIONS ARE MADE WITHOUT GUARANTEE, EXPRESS OR IMPLIED, ON OUR PART. WE DISCLAIM ALL LIABILITY IN CONNECTION WITH THE USE OF THE INFORMATION CONTAINED HEREIN OR OTHERWISE, AND
ALL SUCH RISKS ARE ASSUMED BY THE USER. WE FURTHER EXPRESSLY DISCLAIM ALL WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE. V1-09809

- 5. Add 5 ml of Universal Neutralization Buffer (NB001) and spin down the cells with an 800 g centrifugation for 5 mins.
- 6. Re-suspend the cell pellet with 10 ml or 15 ml of Universal Full Growth Medium and transfer 5 ml each into 2 or 3 pre-coated T25 flasks (for 1/2 to 1/3 subculture ratio).
- 7. Change the medium every 2 or 3 days and the cells usually become confluent within 7 days (when split at a 1/3 ratio).

## FOR RESEARCH USE ONLY

NEUROMICS REAGENTS ARE FOR IN VITRO AND CERTAIN NON-HUMAN IN VIVO EXPERIMENTAL USE ONLY AND NOT INTENDED FOR USE IN ANY HUMAN CLINICAL INVESTIGATION, DIAGNOSIS, PROGNOSIS, OR TREATMENT. THE ABOVE ANALYSES ARE MERELY TYPICAL GUIDES. THEY ARE NOT TO BE CONSTRUED AS BEING SPECIFICATIONS. ALL OF THE ABOVE INFORMATION IS, TO THE BEST OF OUR KNOWLEDGE, TRUE AND ACCURATE. HOWEVER, SINCE THE CONDITIONS OF USE ARE BEYOND OUR CONTROL, ALL RECOMMENDATIONS OR SUGGESTIONS ARE MADE WITHOUT GUARANTEE, EXPRESS OR IMPLIED, ON OUR PART. WE DISCLAIM ALL LIABILITY IN CONNECTION WITH THE USE OF THE INFORMATION CONTAINED HEREIN OR OTHERWISE, AND ALL SUCH RISKS ARE ASSUMED BY THE USER. WE FURTHER EXPRESSLY DISCLAIM ALL WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE, V1-09809